



This plan should be completed by the student's personal diabetes health care team, including the parents/guardian. It should be reviewed with relevant school staff and copies should be kept in a place that can be accessed easily by the school nurse, trained diabetes personnel, and other authorized personnel.

Date of Plan:	_ This plan is va	alid for the current school year:	
Student's Name:		Date of Birth:	
Date of Diabetes Diagnosis:		type 1 type 2 Other	
School:	School	ol Phone Number:	
	Homeroom Tea		
School Nurse:		Phone:	
CONTACT INFORMATION			
Mother/Guardian:	_ Address:		
Telephone			
		Cell:	
Address:			
Telephone: Home	Work	Cell:	
Email Address:			
Student's Physician/Health Ca	are Provider:		
Address:			
Telephone:			
Email Address:	Emerg	gency Number:	
Other Emergency Contacts:			
Name:	Relat	ionship:	

Telephone: Home		Work		Cell:	
-----------------	--	------	--	-------	--

CHECKING BLOOD GLUCOSE

Target range of blood glucose:70-130 mg/dL70-180 mg/dL
Other:
Check blood glucose level: Before lunch Hours after lunch
☐ 2 hours after a correction dose ☐ Mid-morning ☐ Before PE ☐ After PE
Before dismissal Other:
As needed for signs/symptoms of low or high blood glucoseAs needed for signs/symptoms of illness
Preferred site of testing:
Brand/Model of blood glucose meter:
Note: The fingertip should always be used to check blood glucose level if hypoglycemia is suspected.
Student's self-care blood glucose checking skills:
☐ Independently checks own blood glucose
May check blood glucose with supervision
Requires school nurse or trained diabetes personnel to check blood glucose
Continuous Glucose Monitor (CGM): Yes No Brand/Model: Alarms set for: (low) and (high)
Note: Confirm CGM results with blood glucose meter check before taking action on sensor blood glucose level. If student has symptoms or signs of hypoglycemia, check fingertip blood glucose level regardless of CGM
HYPOGLYCEMIA TREATMENT
Student's usual symptoms of hypoglycemia (list below):
If exhibiting symptoms of hypoglycemia, OR if blood glucose level is less thanmg/dL, give a quick-acting glucose product equal to grams of carbohydrate.
Recheck blood glucose in 10-15 minutes and repeat treatment if blood glucose level is less thanmg/dL.

Additional treatment:

HYPOGLYCEMIA TREATMENT (Continued)

Follow physical activity and sports orders (see page 7).
 If the student is unable to eat or drink, is unconscious or unresponsive, or is having seizure activity or convulsions (jerking movements), give: Glucagon: 1 mg 1/2 mg Route: SC IM
• Site for glucagon injection: arm thigh Other:
• Call 911 (Emergency Medical Services) and the student's parents/guardian.
• Contact student's health care provider.
HYPERGLYCEMIA TREATMENT Student's usual symptoms of hyperglycemia (list below):
Check Urine Blood for ketones every hours when blood glucose levels
are above mg/dL.
For blood glucose greater than mg/dL AND at least hours since last insulin dose, give correction dose of insulin (see orders below).
For insulin pump users: see additional information for student with insulin pump.
Give extra water and/or non-sugar-containing drinks (not fruit juices):ounces per

Follow physical activity and sports

Additional treatment for ketones:

orders (see page 7).

hour.

- Notify parents/guardian of onset of hyperglycemia.
- If the student has symptoms of a hyperglycemia emergency, including dry mouth, extreme thirst, nausea and vomiting, severe abdominal pain, heavy breathing or shortness of breath, chest pain, increasing sleepiness or lethargy, or depressed level of consciousness: Call 911 (Emergency Medical Services) and the student's parents/guardian.
- Contact student's health care provider.

Diabetes Medical Management Plan (DMMP) — page 4
INSULIN THERAPY
Insulin delivery device: syringe insulin pen insulin pump
Type of insulin therapy at school: Adjustable Insulin Therapy Fixed Insulin Therapy No insulin
Adjustable Insulin Therapy
• Carbohydrate Coverage/Correction Dose:
Name of insulin:
• Carbohydrate
Coverage:
Insulin-to-Carbohydrate Ratio:
Lunch: 1 unit of insulin per grams of carbohydrate
Snack: 1 unit of insulin per grams of carbohydrate
Carbohydrate Dose Calculation Example Grams of carbohydrate in meal Insulin-to-carbohydrate ratio = units of insulin
• Correction Dose:
<u>Correction Dose Calculation Example</u> <u>Actual Blood Glucose—Target Blood Glucose</u> <u>Blood Glucose Correction Factor/Insulin Sensitivity Factor</u> =units of insulin
Blood Glucose Correction Factor/Insulin Sensitivity Factor =
Target blood glucose = mg/dL
Correction dose scale (use instead of calculation above to determine insulin correction dose):
Blood glucose to mg/dL give units
Blood glucose tomg/dL give units
Blood glucose to mg/dL give units
Blood glucose to mg/dL give units

INSULIN THERAPY (Continued)

When to give insu	ılin:
Lunch Carbohydrate of	coverage only
mg/dL a	coverage plus correction dose when blood glucose is greater than andhours since last insulin dose.
Other:	
Snack No coverage for	or snack
Carbohydrate o	coverage only
Carbohydrate c	overage plus correction dose when blood glucose is greater than
	and hours since last insulin dose.
Correction dose	e only:
For blood glucose ginsulin dose.	greater thanmg/dL AND at least hours since last
Other:	
Fixed Insulin Ther	ару
Name of insulin:	
Units of	
insulin given pre-lu	ınch daily
Units of i	insulin given pre-snack daily
Other:	
Parental Authoriz	ation to Adjust Insulin Dose:
Yes No	Parents/guardian authorization should be obtained before administering a correction dose.
Yes No	Parents/guardian are authorized to increase or decrease correction dose scale within the following range: +/units of insulin.
Yes No	Parents/guardian are authorized to increase or decrease insulin-to-carbohydrate ratio within the following range: units per prescribed grams of carbohydrate, +/ grams of carbohydrate.
Yes No	Parents/guardian are authorized to increase or decrease fixed insulin dose within the following range: +/ units of insulin.

INSULIN THERAPY (Continued)

Student's self-care insulin administration skill					
Yes No Independently calculates and gives	own injections				
Yes No May calculate/give own injections with supervision					
Yes No Requires school nurse or trained di injections	abetes personnel to calculate/give				
ADDITIONAL INFORMATION FOR STUDENT	WITH INSULIN PUMP				
Brand/Model of pump: Type	e of insulin in pump:				
Basal rates during school:					
Type of infusion set:					
For blood glucose greater thanmg/dL =hours after correction, consider pump f parents/guardian.					
☐ For infusion site failure: Insert new infusion set a	nd/or replace reservoir.				
For suspected pump failure: suspend or remove p pen.	oump and give insulin by syringe or				
Physical Activity					
May disconnect from pump for sports activities	Yes No				
Set a temporary basal rate Yes No——— Suspend pump use Yes No	% temporary basal for hours				
Student's self-care pump skills:	Independent?				
Count carbohydrates	Yes No				
Bolus correct amount for carbohydrates consumed	☐ Yes ☐ No				
Calculate and administer correction bolus	☐ Yes ☐ No				
Calculate and set basal profiles	☐ Yes ☐ No				
Calculate and set temporary basal rate	Yes No				
Change batteries	☐ Yes ☐ No				
Disconnect pump	Yes No				
Reconnect pump to infusion set	☐ Yes ☐ No				
Prepare reservoir and tubing	☐ Yes ☐ No				
Insert infusion set	Yes No				
Troubleshoot alarms and malfunctions	☐ Yes ☐ No				

OTHER DIABETES ME	DICATIONS		
Name:	Dose: _	Route:	Times given:
Name:	Dose: _	Route:	Times given:
MEAL PLAN			
Meal/Snack	Time	Carbohydrate Conte	ant (arame)
		to	,
_		to	
Mid-afternoon snack			
Other times to give snacks	s and content/amo	ount:	
Instructions for when food sampling event):	-	ne class (e.g., as part of a	class party or food
Special event/party food p	permitted: Par	rents/guardian discretion	
	☐ Stu	ident discretion	
Student's self-care nutr Yes No Indepen		rbohydrates	
☐ Yes ☐ No May co	unt carbohydrate	s with supervision	
Yes No Require carbohy		ained diabetes personnel	to count
PHYSICAL ACTIVITY	AND SPORTS		
A quick-acting source of giuice must be available at			
Student should eat 15 g	grams 🗌 30 gran	ms of carbohydrate 🔲 o	other
☐ before ☐ every 30	minutes during	after vigorous physic	al activity
other			
If most recent blood gluco physical activity when blo	ose is less than _	mg/dL, student c	
Avoid physical activity w blood ketones are modera	-	e is greater than	mg/dL or if urine/
(Additional information for	or student on insu	llin pump is in the insulir	section on page 6.)

DISASTER PLAN To prepare for an unplanned disaster or emerg supply kit from parent/guardian. Continue to follow orders contained in this Additional insulin orders as follows: Other:	is DMMP.			
SIGNATURES				
This Diabetes Medical Management Plan has	been approved by:			
Student's Physician/Health Care Provider	Date			
I, (parent/guardian:) give permission to the school nurse or another qualified health care professional or trained diabetes personnel of (school:) to perform and carry out the diabetes care tasks as outlined in (student:) 's Diabetes Medical Management Plan. I also consent to the release of the information contained in this Diabetes Medical Management Plan to all school staff members and other adults who have responsibility for my child and who may need to know this information to maintain my child's health and safety. I also give permission to the school nurse or another qualified health care professional to contact my child's physician/health care provider.				
	Date			
Acknowledged and received by:	Date			
Student's Parent/Guardian	Date			

C	tuc	lent	'c I	Dare	nt	/G	119	rd	ia	n
٠,	1111	ш	S F	- 2116	5111	/ L T	па	1(1	14	ш

School Nurse/Other Qualified Health Care Personnel